

# **THE $\alpha_2\delta$ SUBUNIT AND ABSENCE EPILEPSY: BEYOND CALCIUM CHANNELS?**

**Roberta Celli<sup>1\*</sup>, Ines Santolini<sup>1\*</sup>, Michela Guiducci<sup>2</sup>, Gilles Van Luijtelaar<sup>3</sup>, Pasquale Parisi<sup>2</sup>,  
Pasquale Striano<sup>4</sup>, Roberto Gradini<sup>1,5</sup>, Giuseppe Battaglia<sup>1</sup>, Richard T. Ngomba<sup>6</sup>, and  
Ferdinando Nicoletti<sup>1,7</sup>**

**<sup>1</sup> I.R.C.C.S. Neuromed, Neuropharmacology Unit, Parco Tecnologico, Pozzilli, Isernia, Italy;  
Departments of <sup>2</sup>Neurosciences, Mental Health and Sensory Organs, <sup>5</sup>Experimental Medicine,  
and <sup>7</sup>Physiology and Pharmacology, University Sapienza of Rome, Italy; <sup>3</sup>Donders Centre for  
Cognition, Donders Institute for Brain, Cognition and Behaviour, Radboud University,  
Nijmegen, The Netherlands; <sup>4</sup>Pediatric Neurology and Muscular Diseases Unit, Department  
of Neurosciences, Rehabilitation, Ophthalmology, Genetics, Maternal and Child Health,  
University of Genoa, "G. Gaslini" Institute, Genova, Italy; <sup>6</sup>University of Lincoln, School of  
Pharmacy, Brayford Pool, Lincoln, LN6 7TS, UK.**

**\*Co-first Authors**

**Address all correspondence to:**

**Ferdinando Nicoletti**

**Department of Physiology and Pharmacology, University Sapienza of Rome**

**IRCCS Neuromed Pozzilli, Italy**

**Tel. 39 0865 915211**

**Fax 39 0865 927575**

**e-mail: [ferdinandonicoletti@hotmail.com](mailto:ferdinandonicoletti@hotmail.com)**



## ***Abstract***

The spike-wave discharges (SWDs) underlying absence seizures, are generated by hyperactivity of T-type voltage sensitive calcium channels (VSCCs) within the cortico-thalamo-cortical network, and the first-line drug used in the treatment of absence epilepsy inhibit T-type calcium channels. The  $\alpha_2\delta$  subunit associates with all types of VSCCs, including T channels, and facilitates VSCC activation. Hence, one expects that drugs that bind to, and inhibit the  $\alpha_2\delta$  subunit, e.g., gabapentin and pregabalin, are protective against absence epilepsy and that mice lacking the  $\alpha_2\delta$  subunit are resistant to evoked absence seizures. In contrast, gabapentin and pregabalin are not clinically useful, and may even be detrimental, in the treatment of absence epilepsy, and *ducky* mice lacking the  $\alpha_2\delta$  subunit *develop* absence seizures. This suggests that the  $\alpha_2\delta$  subunit displays functions that go beyond the regulation of VSCCs, and that these functions are involved in the regulation of the cortico-thalamo-cortical network. This viewpoint critically examines the role of the  $\alpha_2\delta$  subunit in the pathophysiology of absence seizures focusing on the potential role of the  $\alpha_2\delta$  ligands, thrombospondins.

## ***1. Role for the $\alpha_2\delta$ subunit in the modulation of VSCCs***

VSCCs are formed by a multimolecular complex that includes the  $\alpha_1$ ,  $\beta_1$ ,  $\gamma$ , and  $\alpha_2\delta$  subunits. The  $\alpha_1$  subunit forms the ion channel and accounts for the basic properties of the channel. Based on the specific type of the  $\alpha_1$  subunit, VSCCs are classified into T-type (containing the  $\alpha_1G$ ,  $\alpha_1H$  and  $\alpha_1I$  subunits), L-type (containing the  $\alpha_1C$ ,  $\alpha_1D$ ,  $\alpha_1S$ , and  $\alpha_1F$  subunits), P/Q-type (containing the  $\alpha_1A$  subunit), N-type (containing the  $\alpha_1B$  subunit), and R-type (containing the  $\alpha_1E$  subunit). There are four isoforms of the intracellular  $\beta$  subunit, and eight isoforms of the transmembrane  $\gamma$  subunits, which do not confer a particular specificity to the VSCCs, although the  $\gamma$  subunit is preferentially found in L-type channels of the skeletal muscle. The  $\gamma_2$  subunit belongs to the group of transmembrane AMPA-receptor regulating proteins (TARPs) and is also named stargazine (reviewed by Tomita et al., 2007; Kato et al., 2010). Interestingly, *stargazer* mice lacking the  $\gamma_2$  subunit show a severe neurological phenotype characterized by ataxia and absence epilepsy (Chen et al., 2000; Coombs and Cull-Candy, 2009; Menuz and Nicoll, 2008; Payne, 2008). Convincing evidence for the role of this protein in absence epilepsy comes from a study in adult GAERS (Genetic Absence Epilepsy Rats of Strasbourg), a well validated genetic absence model (Depaulis et al., 2016), in which it was found that elevated levels of stargazin in the cortical focal region (the somatosensory cortex) are associated with an increase in AMPA receptor proteins, GluA1 and GluA2. Elevated stargazin expression is not seen in the epileptic WAG/Rij rat (Wistar Albino Glaxo rats from Rijswijk), another well described genetic absence model endowed with face, construct, and pharmacological validity (Coenen and van Luijtelaar, 2003; Ngomba et al., 2011), suggesting that these changes are not likely a consequence of the hundreds daily SWDs, but it might be one of the many causal factors that play a role in the occurrence of absence seizures and or in epileptogenesis.

The transmembrane  $\alpha_2\delta$  subunit is formed by the  $\alpha_2$  and  $\delta$  subunits covalently linked by a disulfide bridge. Interestingly, the two subunits are generated by a single gene product, which is post-translationally cleaved. The  $\alpha_2$  subunit is glycosylated and interacts extracellularly with the  $\alpha_1$

subunit. The  $\delta$  subunit has a single transmembrane domain and interacts with anchoring intracellular proteins *via* the C-terminal region (De Jongh et al., 1990). There are four isoforms of the  $\alpha_2\delta$  subunit, produced by different genes.  $\alpha_2\delta$ -1, originally described in the skeletal muscle, is ubiquitous;  $\alpha_2\delta$ -2 and  $\alpha_2\delta$ -3 are specifically found in neurons, whereas  $\alpha_2\delta$ -4 is non-neuronal.

The  $\alpha_2\delta$ -1 protein is mainly present in axon terminals of excitatory neurons (Cole et al., 2005), whereas the  $\alpha_2\delta$ -2 protein is expressed by cerebellar Purkinje cells (Barclay et al., 2001). The  $\alpha_2\delta$ -3 protein is ubiquitously expressed in the forebrain (Cole et al., 2005).

The  $\alpha_2\delta$  subunit regulates the activity of VSCCs by increasing current amplitude and causing a shift of voltage-dependent activation towards more hyperpolarized membrane potentials (Arikkath and Campbell, 2003; Dolphin, 2003; Cantí et al., 2005; Hendrich et al., 2008; Davies et al., 2010). In addition, the  $\alpha_2\delta$  subunit regulates cellular trafficking and membrane expression of VSCCs. Accordingly, transient expression of the  $\alpha_2\delta$ -1, -2, and -3 subunits in cultured hippocampal neurons targets P/Q channels in presynaptic terminals, thereby enhancing depolarization-evoked neurotransmitter release (Hoppa et al., 2012). This mechanism may be particularly relevant under pathological conditions, as suggested by the evidence that the  $\alpha_2\delta$ -1 subunit translocates from neuronal cell bodies of dorsal root ganglia to axon terminals in the superficial layers of the dorsal horns of the spinal cord in a mouse model of neuropathic pain (Bauer et al., 2009).

## ***2. The ambiguous role for the $\alpha_2\delta$ subunit in absence epilepsy***

### **a) T-types VSCCs: the key molecular players in absence epilepsy**

Absence seizures, the hallmark of absence epilepsy, are characterized by transient lapses of consciousness associated with bilateral symmetrical SWDs in the electroencephalogram (EEG). Absence seizures are generated by pathological oscillations in a cortico-thalamo-cortical network with a cortical origin; sleep spindles, another type of thalamo-cortical oscillations, are generated in the reticular thalamic nucleus, which is part of this network. This SWDs generating network includes interconnected pyramidal cells and GABA-ergic interneurons in the somatosensory cortex

(SSCtx), thalamo-cortical relay cells in the ventrobasal thalamic (VBT) nuclei, and the reticular thalamic nucleus (RTN) (reviewed by Blumenfeld, 2005; Avoli and Gloor, 1982; Meeren et al., 2005; van Luijtelaar et al., 2011). Highly excitable cells in deep layers of the somatosensory cortex send excitatory fibers to both the VBT and RTN, VBT neurons in their turn send excitatory fibers to the cortical pyramidal neurons and to the GABA-ergic RTN. RTN neurons project to the VBT, and are interconnected by gap junction (reviewed by Blumenfeld, 2005; Ngomba et al., 2011; Kohmann et al., 2016).

T-type VSCCs play a critical role in the generation of pathological oscillations underlying the electrophysiological correlate of absence seizures, the SWDs. These channels are activated at negative membrane potentials by the hyperpolarization-activated cyclic nucleotide-gated cationic  $I_h$  pacemaker channels (HCN), and show a fast voltage-dependent inactivation with respect to other VSCCs. Owing to these peculiar functional properties, T channels are involved in the generation of repetitive firing in the thalamo-cortical network (Perez-Reyes, 2006; Talavera and Nilius, 2006). In the generation of SWDs, an increased excitatory drive from the somatosensory cortex to the RTN generates pathological bursts of RTN GABAergic projection neurons leading to GABA<sub>B</sub> receptor-mediated inhibitory postsynaptic potentials (IPSPs) in VBT neurons. The resulting hyperpolarization enhances the activity of T-type VSCCs in VBT neurons producing the pathological 3-4 Hz oscillations that are typical of absence seizures in humans and 7-11 Hz in the genetic rat models (reviewed by Blumenfeld, 2005). Based on pioneering in vitro studies it is thought that the therapeutic activity of ethosuximide in absence epilepsy occurs via reducing the low-threshold  $Ca^{2+}$  current ( $I_T$ ) in thalamocortical neurons (Coulter et al., 1989; Crunelli and Leresche, 2002). Although modern views emphasize a leading role for the cortex in the initiation of SWDs, the above described thalamic cellular properties, the anatomical projection and an intact cortico-thalamo-cortical network remain imperative for SWDs to occur.

There are three subtypes of T-type channels: Cav3.1, Cav3.2 and Cav3.3, containing the pore-forming  $\alpha_1G$ ,  $\alpha_1H$  and  $\alpha_1I$  subunits, respectively (Catterall et al., 2005). Cav3.2 and Cav3.3 channels

are expressed in RTN neurons and at least Cav3.3 has been implicated in the generation of RTN bursts. Cav3.1 channels are highly expressed in thalamocortical neurons, and all types of T channels are found in cortical neurons (Ernst et al., 2009; Cain and Snutch, 2010; Cheong and Shin, 2013).

#### **b) The $\alpha_2\delta$ subunit associates with T channels and enhances T channel activity in heterologous expression systems**

A few studies suggest that the  $\alpha_2\delta$  subunit positively modulates the activity of T channels, although the physiological significance of the association between the  $\alpha_2\delta$  subunit and T channels is still debated. Overexpression of  $\alpha_2\delta$  influences the voltage-dependence of activation of native T-type channels in neuroblastoma/glioma cell lines (Wyatt et al., 1998). In *Xenopus* oocytes and mammalian cells the  $\alpha_2\delta$  subunit enhances membrane expression and current amplitude of T channels (Dolphin et al., 1999; Dubel et al., 2004; Thompson et al., 2011). The  $\alpha_2\delta$  subunit can also alter the activation and inactivation kinetics and gating properties of T channels (Hobom et al., 2000; Lacinová and Klugbauer, 2004).

#### **c) The *ducky* mouse: a counterintuitive association between the lack of $\alpha_2\delta$ subunit and absence epilepsy**

The association between the *Cacna2d2* gene and childhood absence epilepsy suggests a role for the  $\alpha_2\delta$  subunit in the pathophysiology of absence seizures (Chioza et al., 2009). Because the  $\alpha_2\delta$  subunit facilitates the activity of T channels (see above) it is reasonable to predict that the lack of the  $\alpha_2\delta$  subunit confers protection against absence seizures. In contrast, *ducky* mice, characterized by a spontaneous loss-of-function mutation of the *Cacna2d2* gene, show generalized bilateral SWDs with a frequency of 5-7 Hz, and are considered as a genetic model for absence epilepsy (Barclay et al., 2001). *Ducky* mice also show severe ataxia, dyskinesias, dysgenesis of the cerebellum and other central nervous system (CNS) regions, axonal dystrophy and demyelination

(Meier, 1968). Another spontaneous mutation of the *Cacna2d2* gene (*entla*) encodes a mutant form of the  $\alpha_2\delta$ -2 protein with an intact C-terminus. This mutation also causes generalized epilepsy (Brill et al., 2004), as does a targeted knockout of *Cacna2d2* (Ivanov et al., 2004). The epileptic phenotype of mice lacking  $\alpha_2\delta$ -2 is counterintuitive if one postulates that the  $\alpha_2\delta$  subunit positively modulates T-type channels in thalamic neurons.

#### **d) Lack of therapeutic activity of the two $\alpha_2\delta$ ligands, gabapentin and pregabalin, in absence epilepsy**

Gabapentin and pregabalin have been developed for the treatment of focal epilepsy and are currently used in the treatment of neuropathic pain, migraine, anxiety disorders, and bipolar disorders (Bialer, 2012; Johannessen Landmark, 2008). Gabapentin and pregabalin bind selectively to  $\alpha_2\delta$ -1 and  $\alpha_2\delta$ -2 with nanomolar affinity (reviewed by Cantí et al., 2003; Klugbauer et al., 2003; Davies et al., 2007; Lana et al., 2014). An arginine residue in position 217 of  $\alpha_2\delta$ -1 is part of the binding pocket, and binding of both [ $^3\text{H}$ ]gabapentin and [ $^3\text{H}$ ]pregabalin is largely reduced in the brain regions of transgenic mice with the  $\alpha_2\delta$ -1 R217A mutation (Gee et al., 1996; Gong et al., 2001; Hendrich et al., 2008; reviewed by Dooley et al., 2007; Dolphin, 2012). Interaction of gabapentin and pregabalin with  $\alpha_2\delta$  has two major effect on VSCCs: (i) a reduced membrane localization of  $\alpha_2\delta$  and  $\alpha_1$  subunits; and (ii) a reduced calcium channel current. It is generally believed that both effects contribute to the therapeutic action of gabapentin and pregabalin in focal epilepsy, neuropathic pain, and generalized anxiety disorder (Dooley et al., 2007, Stahl et al., 2013). By interacting with the  $\alpha_2\delta$  subunit, gabapentin and pregabalin are expected to have therapeutic potential in absence epilepsy as a result of T channel inhibition. In contrast, clinical studies have consistently shown that the two drugs have no activity in absence epilepsy (Trudeau et al., 1996; Chadwick et al., 1996), and gabapentin can precipitate absence and myoclonic status epilepticus (Perrucca et al., 1998; Thomas et al., 2006). In animal studies, pregabalin did not affect the



incidence of SWDs in GAERS (Vartanian et al., 2006) and gabapentin had no effect on absence-like seizures in the lethargic mouse model of human absence epilepsy (Hosford and Wang, 1997).

### ***3. Potential explanations for the ambiguous role of the $\alpha_2\delta$ subunit in absence epilepsy***

#### **a) Modulation of non-T type VSCCs**

One possible explanation for the epileptic phenotype of *ducky* and *entla* mice is that the loss of  $\alpha_2\delta$  subunit causes absence seizures by restraining the activity of high voltage-activated (HVA)  $\text{Ca}^{2+}$  channels (P/Q, N, L and R channels). Although T channels are one of the major players in the generation of pathological oscillations in the cortico-thalamo-cortical network, a large body of evidence suggests that HVA channels are also involved in the pathophysiology of absence seizures. Pharmacological activation of L-type HVA  $\text{Ca}^{2+}$  channels alters the firing rate of thalamocortical neurons (Kanyshkova et al., 2014), while the systemic administration of L-type  $\text{Ca}^{2+}$  channel blocker nimodipine increased SWDs, while a decrease was found after i.c.v. (intracerebroventricular) BAY K8644, a L-type  $\text{Ca}^{2+}$  channel opener. BAY K8644 was also able to antagonise the effects of nimodipine in rats with spontaneous occurring SWDs (van Luijtelaar et al., 1995; van Luijtelaar et al., 2000). R-type channels are also involved in the generation of oscillatory burst discharges in RTN neurons and absence epilepsy (Zaman et al., 2011). Preliminary evidence for a role in the R-type  $\text{Ca}^{2+}$  blocker was obtained after the administration of the conotoxin GVIA, it reduced SWDs (van Luijtelaar et al., 2000). Mutations of the *Cacna1a* gene encoding for the  $\alpha_1$  subunit of P-type channels are associated with absence seizures in rodents and humans (Burgess and Noebels, 1999; Zamponi et al., 2010; Serikawa et al., 2015). It is possible that in *ducky* mice a reduced expression/activity of HVA VSCCs causes a compensatory increase in the activity of T-type channels resulting into absence seizures.

#### **b) Beyond VSCCs: the $\alpha_2\delta$ subunit mediates the action of thrombospondins in the CNS**

The  $\alpha_2\delta$  subunit is the receptor for a class of proteins, called thrombospondins (TSPs), which were first isolated from platelets stimulated with thrombin and show a widespread distribution in various organs, including the CNS. TSPs are a large oligomeric extracellular matrix (ECM) proteins, found in vertebrates and lower metazoa (Adams, 2001; Bentley and Adams, 2010; Mosher and Adams, 2012). The human TSP protein family consists of five members (TSP-1, TSP-2, TSP-3, TSP-4, and TSP-5, or COMP, cartilage oligomeric matrix protein) which are subdivided into two groups (Adams, 2001; Adams and Lawler, 2004; Adams and Lawler, 2011) on the basis of their oligomerization state and domain structure. Subgroup A includes TSP-1 and TSP-2, which form homotrimers; subgroup B includes TSP-3, TSP-4, and TSP-5, which form homopentamers (Lawler, 2002). The various isoforms of TSPs are the products of different genes, and display differential non-overlapping functions (Stenina-Adognravi, 2013).

TSPs primarily regulate cell-cell and cell-extracellular matrix interactions (Bornstein, 2000) acting through a number of ECM proteins and cell surface receptors and controlling cytoskeletal dynamics, cell migration, and cell attachment (Bornstein et al., 2004). All these activities regulate many aspects of cell phenotype and contribute to angiogenesis and wound healing, vessel wall biology and synaptogenesis (Adams and Lawler, 2011).

Astrocyte-secreted TSP-1, TSP-2 promote synaptogenesis in cultured hippocampal neurons (Xu et al., 2010) and retinal ganglion cells (Christopherson et al., 2015), and TSP-1 can also promote neuronal migration (Iruela-Arispe et al., 1993) and axonal growth (Yu et al., 2008). TSP-1 and TSP-2 are highly expressed during development (Christopherson et al., 2005; Xu et al., 2010; Cahoy et al., 2008) and acts in specific time windows to promote synaptogenesis in the developing brain. TSP-4 is exclusively expressed by mature astrocytes, and represents the mature TSP isoform in the CNS (Bornstein et al., 2004).

Eroglu et al. (2009) have shown for the first time that the synaptogenic epidermal growth factor (EGF)-like repeat domains of TSPs interact with the VWF-A (von Willebrand factor) domain of  $\alpha_2\delta$ -1. This interaction mediates the synaptogenic activity of TSPs. Accordingly, overexpression of

$\alpha_2\delta$ -1 enhanced synaptogenesis in mice, and the  $\alpha_2\delta$  ligand, gabapentin, restrained the synaptogenic activity of TSPs (Eroglu et al., 2009).

A potential link between alterations in synaptogenesis and absence epilepsy is suggested by the evidence that stargazin mutation in mice causes an impairment in synaptic formation and maturation in the cerebellum (Meng et al., 2006). In addition, double mutant *zi/zi*, *tm/tm* rats, which spontaneously develop absence seizures after 6 weeks of age, show a low expression of the synaptic vesicle proteins, SV2A and synaptotagmin-1 (Hanaya et al., 2012). We have found a selective reduction of the transcript encoding for TSP-1 in the ventrobasal thalamus of WAG/Rij (Santolini et al., 2015), which represent a validated genetic rat model of human absence epilepsy (van Luijtelaar and Coenen, 1986; 1997).

An attractive hypothesis is that the TSP- $\alpha_2\delta$  axis plays a key role in synaptic formation within the cortico-thalamo-cortical network that lies at the core of the pathophysiology of absence epilepsy (Blumenfeld, 2005) and that defects in either TSPs or  $\alpha_2\delta$  disrupt synaptic organization in the circuit resulting into absence seizures. TSPs may have a broader role in epileptogenesis that involves other types of epilepsy. Mendus et al. (2015) have recently found that mice with genetic deletion of TSP-1 show an increased sensitivity to chemical kindling induced by repeated administrations of a subconvulsive dose of pentylenetetrazole (PTZ). TSP-1 knockout mice developed generalized tonic-clonic seizures in response to the second administration of PTZ, whereas at least eight administrations were required to produce the same phenotype in wild-type mice (Mendus et al., 2015). Interestingly, TSP-1 knockout mice also showed a reduced expression of both  $\alpha_2\delta$ -1 and  $\alpha_2\delta$ -2 subunits in the frontal cortex (Mendus et al., 2015).

In conclusion, the  $\alpha_2\delta$  subunit subserves pleiotropic functions that may be relevant for the pathophysiology of epilepsy and may be targeted by therapeutic intervention. The role of  $\alpha_2\delta$  subunit in absence epilepsy cannot be reconducted to the modulation of T-type channels, and may involve other types of VSCCs. Increasing evidence suggests that the TSP- $\alpha_2\delta$  axis can be dysfunctional in absence epilepsy and other forms of epilepsy. Abnormalities in either TSPs and/or

$\alpha_2\delta$  subunit may contribute to the process of epileptogenesis, which is ultimately responsible for the recurrence of epileptic seizures. If proven to be correct, this may lay the groundwork for the design of novel disease modifying drugs that are directed towards mechanisms that lie at the core of epileptogenesis.

## REFERENCES

- Adams JC. Thrombospondins: multifunctional regulators of cell interactions. *Annu Rev Cell Dev Biol.* 2001;17:25-51.
- Adams JC, Lawler J. The thrombospondins. *Int J Biochem Cell Biol.* 2004; 36:961–968.
- Adams JC, Lawler J. The thrombospondins. *Cold Spring Harb Perspect Biol.* 2011 Oct1; 3(10):a009712.
- Arikkath J, Campbell KP. Auxiliary subunits: essential components of the voltage-gated calcium channel complex. *Curr Opin Neurobiol.* 2003 Jun;13(3):298-307.
- Avoli M, Gloor P. Interaction of cortex and thalamus in spike and wave discharges of feline generalized penicillin epilepsy. *Exp Neurol.* 1982 Apr;76(1):196-217.
- Barclay J, Balaguero N, Mione M, Ackerman SL, Letts VA, Brodbeck J, Canti C, Meir A, Page KM, Kusumi K, Perez-Reyes E, Lander ES, Frankel WN, Gardiner RM, Dolphin AC, Rees M. Ducky mouse phenotype of epilepsy and ataxia is associated with mutations in the *Cacna2d2* gene and decreased calcium channel current in cerebellar Purkinje cells. *J Neurosci.* 2001 Aug 15;21(16):6095-104.
- Bauer CS, Nieto-Rostro M, Rahman W, Tran-Van-Minh A, Ferron L, Douglas L, Kadurin I, Sri Ranjan Y, Fernandez-Alacid L, Millar NS, Dickenson AH, Lujan R, Dolphin AC. The increased trafficking of the calcium channel subunit  $\alpha 2\delta$ -1 to presynaptic terminals in neuropathic pain is inhibited by the  $\alpha 2\delta$  ligand pregabalin. *J Neurosci.* 2009 Apr 1;29(13):4076-88.
- Bentley AA, Adams JC. The evolution of thrombospondins and their ligand-binding activities. *Mol Biol Evol.* 2010 Sep;27(9):2187-97.
- Bialer M. Why are antiepileptic drugs used for nonepileptic conditions? *Epilepsia.* 2012 Dec;53 Suppl 7:26-33.

Blumenfeld H. Cellular and network mechanisms of spike-wave seizures. *Epilepsia*. 2005;46 Suppl 9:21-33.

Bornstein P. Matricellular proteins: an overview. *Matrix Biol*. 2000 Dec;19(7):555-6.

Bornstein P, Agah A, Kyriakides TR. The role of thrombospondins 1 and 2 in the regulation of cellmatrix interactions, collagen fibril formation, and the response to injury. *Int J Biochem Cell Biol*. 2004; 36:1115–25.

Brill J, Klocke R, Paul D, Boison D, Gouder N, Klugbauer N, Hofmann F, Becker CM, Becker K. entla, a novel epileptic and ataxic *Cacna2d2* mutant of the mouse. *Journal of Biological Chemistry* 2004;279:7322–7330.

Burgess DL, Noebels JL. Single gene defects in mice: the role of voltage-dependent calcium channels in absence models. *Epilepsy Res*. 1999 Sep;36(2-3):111-22.

Cahoy JD, Emery B, Kaushal A, et al. A transcriptome database for astrocytes, neurons, and oligodendrocytes: a new resource for understanding brain development and function. *J Neurosci*. 2008;28(1):264–278.

Cain SM, Snutch TP. “Contributions of T-type calcium channel isoforms to neuronal firing.” *Channels (Austin)* (2010)4:475–82.

Cantí C, Davies A, DolphinAC. Calcium Channel  $\alpha_2\delta$  Subunits: Structure, Functions and Target Site for Drugs. *Curr. Neuropharmacol*. 2003; 1(3): 209-217.

Cantí C, Nieto-Rostro M, Foucault I, Heblich F, Wratten J, Richards MW, Hendrich J, Douglas L, Page KM, Davies A, Dolphin AC. The metal-ion-dependent adhesion site in the Von Willebrand factor-A domain of  $\alpha_2\delta$  subunits is key to trafficking voltage-gated  $\text{Ca}^{2+}$  channels. *Proc Natl Acad Sci U S A*. 2005 Aug 9;102(32):11230-5.

Catterall WA, Perez-Reyes E, Snutch TP, Striessnig J. International Union of Pharmacology. XLVIII. Nomenclature and structure-function relationships of voltage-gated calcium channels. *Pharmacol Rev.* 2005 Dec;57(4):411-25.

Chadwick D, Leiderman DB, Sauermann W, Alexander J, Garofalo E. Gabapentin in generalized seizures. *Epilepsy Res.* 1996 Nov;25(3):191-7.

Chen L, Chetkovich DM, Petralia RS, Sweeney NT, Kawasaki Y, Wenthold RJ, Brecht DS, Nicoll RA. Stargazin regulates synaptic targeting of AMPA receptors by two distinct mechanisms. *Nature.* 2000 Dec 21-28;408(6815):936-43.

Cheong E, Shin HS. T-type  $\text{Ca}^{2+}$  channels in absence epilepsy. *Biochim Biophys Acta.* 2013 Jul;1828(7):1560-71.

Chioza BA, Aicardi J, Aschauer H, Brouwer O, Callenbach P, Covanis A, Dooley JM, Dulac O, Durner M, Eeg-Olofsson O, et al. Genome wide high density SNP-based linkage analysis of childhood absence epilepsy identifies a susceptibility locus on chromosome 3p23-p14. *Epilepsy research.* 2009; 87:247–255.

Christopherson KS, Ullian EM, Stokes CC, et al. Thrombospondins are astrocyte-secreted proteins that promote CNS synaptogenesis. *Cell.* 2005;120(3):421–433.

Coenen AM, Van Luijtelaar EL. Genetic animal models for absence epilepsy: a review of the WAG/Rij strain of rats. *Behav Genet.* 2003 Nov;33(6):635-55.

Cole RL, Lechner SM, Williams ME, Prodanovich P, Bleicher L, Varney MA, Gu G. Differential distribution of voltage-gated calcium channel alpha-2 delta (alpha2delta) subunit mRNA-containing cells in the rat central nervous system and the dorsal root ganglia. *J Comp Neurol.* 2005 Oct 24;491(3):246-69.

Coombs ID, Cull-Candy SG. Transmembrane AMPA receptor regulatory proteins and AMPA receptor function in the cerebellum. *Neuroscience*. 2009 Sep 1;162(3):656-65.

Coulter DA, Huguenard JR, Prince DA. Characterization of ethosuximide reduction of low-threshold calcium current in thalamic neurons. *Ann Neurol*. 1989 Jun;25(6):582-93.

Crunelli V, Leresche N. Block of Thalamic T-Type  $\text{Ca}(2+)$  Channels by Ethosuximide Is Not the Whole Story. *Epilepsy Curr*. 2002 Mar;2(2):53-56.

Davies A, Hendrich J, Van Minh AT, Wratten J, Douglas L, Dolphin AC. Functional biology of the  $\alpha(2)\delta$  subunits of voltage-gated calcium channels. *Trends Pharmacol Sci*. 2007 May;28(5):220-8.

Davies A, Kadurin I, Alvarez-Laviada A, Douglas L, Nieto-Rostro M, Bauer CS, Pratt WS, Dolphin AC. The  $\alpha 2\delta$  subunits of voltage-gated calcium channels form GPI-anchored proteins, a posttranslational modification essential for function. *Proc Natl Acad Sci U S A*. 2010 Jan 26;107(4):1654-9.

De Jongh KS, Warner C, Catterall WA. Subunits of purified calcium channels. Alpha 2 and delta are encoded by the same gene. *J Biol Chem*. 1990 Sep 5;265(25):14738-41.

Depaulis A, David O, Charpier S. The genetic absence epilepsy rat from Strasbourg as a model to decipher the neuronal and network mechanisms of generalized idiopathic epilepsies. *J Neurosci Methods*. 2016 Feb 15;260:159-74.

Dolphin AC. G protein modulation of voltage-gated calcium channels. *Pharmacol Rev*. 2003 Dec;55(4):607-27.

Dolphin AC. Calcium channel auxiliary  $\alpha 2\delta$  and  $\beta$  subunits: trafficking and one step beyond. *Nat Rev Neurosci*. 2012 Jul 18;13(8):542-55.



Dolphin AC, Wyatt CN, Richards J, et al. The effect of alpha2-delta and other accessory subunits on expression and properties of the calcium channel alpha1G. *J Physiol*. 1999;519(Pt 1):35–45.

Dooley DJ, Taylor CP, Donevan S, Feltner D. Ca<sup>2+</sup> channel alpha2delta ligands: novel modulators of neurotransmission. *Trends Pharmacol Sci*. 2007 Feb;28(2):75-82.

Dubel SJ, Altier C, Chaumont S, Lory P, Bourinet E, Nargeot J. Plasma membrane expression of T-type calcium channel alpha(1) subunits is modulated by high voltage-activated auxiliary subunits. *J Biol Chem*. 2004 Jul 9;279(28):29263-9.

Ernst WL, Zhang Y, Yoo JW, Ernst SJ, Noebels JL. Genetic enhancement of thalamocortical network activity by elevating alpha 1g-mediated low-voltage-activated calcium current induces pure absence epilepsy. *J Neurosci*. 2009 Feb 11;29(6):1615-25.

Eroglu C, Allen NJ, Susman MW, et al. Gabapentin receptor alpha2delta-1 is a neuronal thrombospondin receptor responsible for excitatory CNS synaptogenesis. *Cell*. 2009;139(2):380–392.

Gee NS, Brown JP, Dissanayake VU, Offord J, Thurlow R, Woodruff GN. The novel anticonvulsant drug, gabapentin (Neurontin), binds to the alpha2delta subunit of a calcium channel. *J Biol Chem*. 1996 Mar 8;271(10):5768-76.

Gong HC, Hang J, Kohler W, Li L, Su TZ. Tissue-specific expression and gabapentin-binding properties of calcium channel alpha2delta subunit subtypes. *J Membr Biol*. 2001 Nov 1;184(1):35-43.

Hanaya R, Hosoyama H, Sugata S, Tokudome M, Hirano H, Tokimura H, Kurisu K, Serikawa T, Sasa M, Arita K. Low distribution of synaptic vesicle protein 2A and synaptotagmin-1 in the cerebral cortex and hippocampus of spontaneously epileptic rats exhibiting both tonic convulsion and absence seizure. *Neuroscience*. 2012 Sep 27;221:12-20.

Hendrich J, Van Minh AT, Hebllich F, Nieto-Rostro M, Watschinger K, Striessnig J, Wratten J, Davies A, Dolphin AC. Pharmacological disruption of calcium channel trafficking by the  $\alpha 2\delta$  ligand gabapentin. *Proc Natl Acad Sci U S A*. 2008 Mar 4;105(9):3628-33.

Hobom M, Dai S, Marais E, Lacinova L, Hofmann F, Klugbauer N. Neuronal distribution and functional characterization of the calcium channel  $\alpha 2\delta$ -2 subunit. *Eur J Neurosci*. 2000 Apr;12(4):1217-26.

Hoppa MB, Lana B, Margas W, Dolphin AC, Ryan TA.  $\alpha 2\delta$  expression sets presynaptic calcium channel abundance and release probability. *Nature*. 2012 May 13;486(7401):122-5.

Hosford DA, Wang Y. Utility of the lethargic (lh/lh) mouse model of absence seizures in predicting the effects of lamotrigine, vigabatrin, tiagabine, gabapentin, and topiramate against human absence seizures. *Epilepsia*. 1997 Apr;38(4):408-14.

Iruela-Arispe ML, Liska DJ, Sage EH, Bornstein P. Differential expression of thrombospondin 1, 2, and 3 during murine development. *Dev Dyn*. 1993 May;197(1):40-56.

Ivanov SV, Ward JM, Tessarollo L, McAreavey D, Sachdev V, Fananapazir L, Banks MK, Morris N, Djurickovic D, Devor-Henneman DE, Wei MH, Alvord GW, Gao B, Richardson JA, Minna JD, Rogawski MA, Lerman MI. Cerebellar ataxia, seizures, premature death, and cardiac abnormalities in mice with targeted disruption of the *Cacna2d2* gene. *Am J Pathol* 2004;165:1007–1018.

Johannessen Landmark C. Antiepileptic drugs in non-epilepsy disorders: relations between mechanisms of action and clinical efficacy. *CNS Drugs*. 2008;22(1):27-47.

Kanyshkova T, Ehling P, Cerina M, Meuth P, Zobeiri M, Meuth SG, Pape HC, Budde T. Regionally specific expression of high-voltage-activated calcium channels in thalamic nuclei of epileptic and non-epileptic rats. *Mol Cell Neurosci*. 2014 Jul;61:110-22.

- Kato AS, Gill MB, Yu H, Nisenbaum ES, Brecht DS. TARPs differentially decorate AMPA receptors to specify neuropharmacology. *Trends Neurosci.* 2010 May;33(5):241-8.
- Klugbauer N, Marais E, Hofmann F. Calcium channel  $\alpha_2\delta$  subunits: differential expression, function, and drug binding. *J Bioenerg Biomembr.* 2003 Dec;35(6):639-47.
- Kohmann D, Lüttjohann A, Seidenbecher T, Coulon P, Pape HC. Short-term depression of gap junctional coupling in reticular thalamic neurons of absence epileptic rats. *J Physiol.* 2016 Oct 1;594(19):5695-710.
- Lacinová L, Klugbauer N. Modulation of gating currents of the  $\text{Ca}_v3.1$  calcium channel by  $\alpha_2\delta_2$  and  $\gamma_5$  subunits. *Arch Biochem Biophys.* 2004 May 15;425(2):207-13.
- Lana B, Schlick B, Martin S, Pratt WS, Page KM, Goncalves L, Rahman W, Dickenson AH, Bauer CS, Dolphin AC. Differential upregulation in DRG neurons of an  $\alpha_2\delta-1$  splice variant with a lower affinity for gabapentin after peripheral sensory nerve injury. *Pain.* 2014 Mar;155(3):522-33.
- Lawler J. Thrombospondin-1 as an endogenous inhibitor of angiogenesis and tumor growth. *J. Cell. Mol. Med.* 2002; 6:1–12.
- Meeren H, van Luijckelaar G, Lopes da Silva F, Coenen A. Evolving concepts on the pathophysiology of absence seizures: the cortical focus theory. *Arch Neurol.* 2005 Mar;62(3):371-6.
- Meier H. The neuropathology of ducky, a neurological mutation of the mouse. A pathological and preliminary histochemical study. *Acta Neuropathol.* 1968 Jul 8;11(1):15-28.
- Mendus D, Rankin-Gee EK, Mustapha M, Porter BE. Increased sensitivity to kindling in mice lacking TSP1. *Neuroscience.* 2015 Oct 1;305:302-8.
- Meng H, Walker N, Su Y, Qiao X. Stargazin mutation impairs cerebellar synaptogenesis, synaptic maturation and synaptic protein distribution. *Brain Res.* 2006 Dec 8;1124(1):197-207.

Menuz K, Nicoll RA. Loss of inhibitory neuron AMPA receptors contributes to ataxia and epilepsy in stargazer mice. *J Neurosci*. 2008 Oct 15;28(42):10599-603.

Mosher DF, Adams JC. Adhesion-modulating/matricellular ECM protein families: a structural, functional and evolutionary appraisal. *Matrix Biol*. 2012 Apr;31(3):155-61.

Ngomba RT, Santolini I, Salt TE, Ferraguti F, Battaglia G, Nicoletti F, van Luijtelaar G. Metabotropic glutamate receptors in the thalamocortical network: strategic targets for the treatment of absence epilepsy. *Epilepsia*. 2011 Jul;52(7):1211-22.

Payne HL. The role of transmembrane AMPA receptor regulatory proteins (TARPs) in neurotransmission and receptor trafficking (Review). *Mol Membr Biol*. 2008 May;25(4):353-62.

Perez-Reyes E. Molecular characterization of T-type calcium channels. *Cell Calcium*. 2006 Aug;40(2):89-96.

Perucca E, Gram L, Avanzini G, Dulac O. Antiepileptic drugs as a cause of worsening seizures. *Epilepsia*. 1998 Jan;39(1):5-17.

Santolini I, Guiducci M, Imbriglio T, Celli R, Cannella M, Motolese M, D'Amore V, Gradini R, van Luijtelaar G, Parisi P, Nicoletti F, Striano P, Ngomba RT. Reduced thalamic expression of thrombospondin-1 in a rat model of spontaneous absence epilepsy. *Society for Neuroscience* 2015, 766.14.

Serikawa T, Mashimo T, Kuramoro T, Voigt B, Ohno Y, Sasa M. Advances on genetic rat models of epilepsy. *Exp Anim*. 2015;64(1):1-7.

Stahl SM, Porreca F, Taylor CP, Cheung R, Thorpe AJ, Clair A. The diverse therapeutic actions of pregabalin: is a single mechanism responsible for several pharmacological activities? *Trends Pharmacol Sci*. 2013 Jun;34(6):332-9.

Stenina-Adognravi O. Thrombospondins: old players, new games. *Curr Opin Lipidol*. 2013 Oct;24(5):401-9.

Talavera K, Nilius B. Biophysics and structure-function relationship of T-type Ca<sup>2+</sup> channels. *Cell Calcium*. 2006 Aug;40(2):97-114.

Thomas P, Valton L, Genton P. Absence and myoclonic status epilepticus precipitated by antiepileptic drugs in idiopathic generalized epilepsy. *Brain*. 2006 May;129(Pt 5):1281-92.

Thompson WR, Majid AS, Czymmek KJ, Ruff AL, García J, Duncan RL, Farach-Carson MC. Association of the  $\alpha(2)\delta(1)$  subunit with Ca(v)3.2 enhances membrane expression and regulates mechanically induced ATP release in MLO-Y4 osteocytes. *J Bone Miner Res*. 2011 Sep;26(9):2125-39.

Tomita S, Shenoy A, Fukata Y, Nicoll RA, Brecht DS. Stargazin interacts functionally with the AMPA receptor glutamate-binding module. *Neuropharmacology*. 2007 Jan;52(1):87-91.

Trudeau V, Myers S, LaMoreaux L, Anhut H, Garofalo E, Ebersole J. Gabapentin in naïve childhood absence epilepsy: results from two double-blind, placebo-controlled, multicenter studies. *J Child Neurol*. 1996 Nov;11(6):470-5.

van Luijtelaar E.L., Ates N., Coenen A.M. Role of L-type calcium channel modulation in nonconvulsive epilepsy in rats. *Epilepsia*, 36 (1995), pp. 86–92.

van Luijtelaar E.L, Coenen A.M. Two types of electrocortical parox-ysms in an inbred strain of rats. *Neurosci Lett* 70 (1986), pp. 393–397.

van Luijtelaar E.L, Coenen A.M. The WAG/Rij rat model of absence epilepsy: ten years of research. Nijmegen University Press, Nijmegen (1997).

van Luijtelaar G, Sitnikova E, Littjohann A. On the origin and suddenness of absences in genetic absence models. *Clin EEG Neurosci.* 2011 Apr;42(2):83-97.

van Luijtelaar G., Wiaderna D., Elants C., Scheenen W. Opposite effects of T- and L-type Ca(2+) channels blockers in generalized absence epilepsy. *Eur. J. Pharmacol.*, 406 (2000), pp. 381–389

Vartanian MG, Radulovic LL, Kinsora JJ, Serpa KA, Vergnes M, Bertram E, Taylor CP. Activity profile of pregabalin in rodent models of epilepsy and ataxia. *Epilepsy Res.* 2006 Mar;68(3):189-205.

Wyatt CN, Page KM, Berrow NS, Brice NL, Dolphin AC. The effect of overexpression of auxiliary Ca<sup>2+</sup> channel subunits on native Ca<sup>2+</sup> channel currents in undifferentiated mammalian NG108-15cells. *J Physiol.* 1998;510 (Pt 2): 347–360.

Xu J, Xiao N, Xia J. Thrombospondin 1 accelerates synaptogenesis in hippocampal neurons through neuroligin 1. *Nat Neurosci.* 2010;13(1):22–24.

Yu K, Ge J, Summers JB, Li F, Liu X, Ma P, Kaminski J, Zhuang J. TSP-1 secreted by bone marrow stromal cells contributes to retinal ganglion cell neurite outgrowth and survival. *PLoS One.* 2008 Jun 25;3(6):e2470.

Zaman T, Lee K, Park C, Paydar A, Choi JH, Cheong E, Lee CJ, Shin HS. Cav2.3 channels are critical for oscillatory burst discharges in the reticular thalamus and absence epilepsy. *Neuron.* 2011 Apr 14;70(1):95-108.

Zamponi GW, Lory P, Perez-Reyes E. Role of voltage-gated calcium channels in epilepsy. *Pflugers Arch.* 2010 Jul;460(2):395-403.



